

# Position of the Working Group Sports Nutrition of the German Nutrition Society (DGE): Energy Needs in Sports

*Position der Arbeitsgruppe Sporternährung der Deutschen Gesellschaft für Ernährung (DGE): Energiebedarf im Sport*

## Summary

- › **Athletes** expend greater amounts of energy as a result of their training. However, these amounts vary during the competition year and are influenced by such factors as the training period and intensity.
- › **Measuring energy expenditure** is a complex process that is often achieved through indirect calorimetry. A calculation is possible when certain limitations are taken into consideration.
- › **Energy expenditure** corresponds to energy requirements. Energy requirements should be covered by energy intake adapted to individual needs. Low energy intake can lead to reduced energy availability. Low energy availability not only has a negative impact on the bone metabolism and menstrual cycle – there may also be consequences of a hematological, metabolic, psychological, gastrointestinal and immunological nature. This has adverse effects on the performance capacity, training adjustment, concentration and coordination, and can lead to an increased risk of injury.
- › **Matching energy intake** to energy expenditure is an important cornerstone in the diet of athletes and a significant prerequisite for maintaining their health and performance.

## KEY WORDS:

Athletes, Calorimetry, Energy Availability, Resting Energy Expenditure, Metabolic Equivalent

## Zusammenfassung

- › **Sportler** haben bedingt durch das Training einen gesteigerten Energieverbrauch. Dieser ist jedoch im Verlaufe eines Wettkampfjahres variabel und wird unter anderem von der Trainingsdauer und -intensität beeinflusst.
- › **Die Messung des Energieverbrauchs** ist komplex und erfolgt häufig über eine indirekte Kalorimetrie. Eine Berechnung ist unter Berücksichtigung bestimmter Einschränkungen möglich.
- › **Der Energieverbrauch** entspricht dem Energiebedarf. Der Energiebedarf sollte durch eine angepasste Energiezufuhr gedeckt werden. Eine geringe Energiezufuhr kann zu einer reduzierten Energieverfügbarkeit führen. Eine niedrige Energieverfügbarkeit hat nicht nur negative Auswirkungen auf den Knochenstoffwechsel und den Menstruationszyklus, sondern es ist auch mit hämatologischen, metabolischen, psychologischen, gastrointestinalen und immunologischen Konsequenzen zu rechnen. Daraus resultieren negative Einflüsse auf die Leistungsfähigkeit, Trainingsanpassung, Konzentration, Koordination und ein erhöhtes Verletzungsrisiko.
- › **Eine dem Energieverbrauch** angepasste Energiezufuhr ist ein wichtiger Eckpfeiler in der Ernährung des Sportlers und eine wesentliche Voraussetzung für Gesundheit und Leistungsfähigkeit.

## SCHLÜSSELWÖRTER:

Athleten, Kalorimetrie, Energieverfügbarkeit, Ruheenergieumsatz, metabolisches Äquivalent

## Introduction

Athletes have special dietary needs. In a consensus statement on sports nutrition, the International Olympic Committee (IOC) stated that diet influences performance capacity and advised athletes to apply dietary strategies adapted to individual needs during and after training and competitions to maximize their physical and mental performance (19). It should be kept in mind that athletes are not a homogeneous group;

there are in fact numerous factors that influence energy and nutrient needs (e.g. body weight, size, body composition, training program, duration, intensity). It must also be considered that, depending on the training and competition phase within the annual cycle and even within any given week, athletes plan the intensity and duration of their training sessions in very different ways (17, 36, 37). ➤

## POSITION STAND

ACCEPTED: August 2020

PUBLISHED ONLINE: September 2020

DOI: 10.5960/dzsm.2020.451

Braun H, Carlsohn A, Großhauser M, König D, Lampen A, Mosler S, Nieß A, Oberritter H, Schäbenthal K, Schek A, Stehle P, Virmani K, Ziegenhagen R, Heseke H. Position of the working group sports nutrition of the German Nutrition Society (DGE): energy needs in sports. Dtsch Z Sportmed. 2020; 71: 171-177.

1. DEUTSCHE SPORHOCHSCHULE KÖLN, Institut für Biochemie; Deutsches Forschungszentrum für Leistungssport, Köln, Germany
2. HOCHSCHULE FÜR ANGEWANDTE WISSENSCHAFTEN HAMBURG, Fakultät Life Sciences/Department Ökotrophologie, Hamburg, Germany
3. OLYMPIASTÜTZPUNKT RHEINLAND-PFALZ/ SAARLAND, Germany
4. ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, Institut für Sport und Sportwissenschaft, Arbeitsbereich Ernährung, Freiburg, Germany
5. BUNDESINSTITUT FÜR RISIKOBEWERTUNG (BfR), Berlin, Germany
6. OLYMPIASTÜTZPUNKT STUTTGART, Germany
7. UNIVERSITÄTSKLINIKUM TÜBINGEN, Abteilung Sportmedizin, Tübingen, Germany
8. DEUTSCHE GESELLSCHAFT FÜR ERNÄHRUNG E. V. (DGE), Germany
9. REDAKTION LEISTUNGSSPORT (DOSB), Germany
10. RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT BONN, Institut für Ernährungs- und Lebensmittelwissenschaften, Ernährungsphysiologie, Bonn, Germany
11. UNIVERSITÄT PADERBORN, Institut für Ernährung, Konsum und Gesundheit, Paderborn, Germany



Article incorporates the Creative Commons Attribution – Non Commercial License. <https://creativecommons.org/licenses/by-nc-sa/4.0/>



Scan QR Code and read article online.

## CORRESPONDING ADDRESS:

Hans Braun  
Deutsche Sporthochschule Köln  
Institut für Biochemie  
Am Sportpark Müngersdorf 6  
50859 Köln, Germany  
✉: h.braun@dshs-koeln.de

Table 1

Selected everyday activities and sports activities with the corresponding METs and codes from the Compendium of Physical Activities (adapted from (5)).

EVERYDAY ACTIVITIES	METS	CODE
Sleeping	0,95	7030
Sitting quietly, e.g. in front of the TV	1,3	7020
Meeting while sitting	1,5	11585
Office work	1,5	11580
Ironing	1,8	5070
Walking leisurely (< 3.2 km/h)	2	11791
Cooking, preparing food	2	5050
Shopping at the supermarket	2,3	5060
Housework, cleaning	2,5	5040
Garden work (leisurely)	3	8260
Garden work (strenuous)	6	8262
SPORTS ACTIVITIES	METS	CODE
Badminton (general)	5,5	15020
Badminton (competitive)	7	15030
Basketball (general)	6	15050
Basketball (competitive)	8	15040
Boxing (with punching bag)	5,5	15110
Boxing (sparring)	7,8	15120
Darts	2,5	15180
Soccer (general)	7	15610
Soccer (competitive)	10	15605
Golf	4,3	15265
Jogging (6.4 km/h)	6	12029
Jogging (8 km/h)	8,3	12030
Jogging (11.2 km/h)	11	12070
Jogging (16 km/h)	14,5	12120
Jogging (22.4 km/h)	23	12135
Biking (9 km/h)	3,5	1018
Biking (18 km/h)	6,8	1020
Biking (24 km/h)	10	1040
Swimming breaststroke (general)	5,3	18255
Swimming breaststroke (intensive)	10,3	18260
Swimming freestyle (general)	5,8	18240
Swimming freestyle (2:10 min/100m)	8,3	18290
Swimming freestyle (intensive)	9,8	18230
Squash (general)	7,3	15652
Tai Chi, Qi Gong	3	15670
Tennis (general)	7,3	15675
Table tennis (general)	4	15660
Volleyball (general)	4	15710
Volleyball (competitive)	6	15711

A young high-performance athlete also has an increased, growth-related need for energy and nutrients (11, 39). Furthermore, the body weight and body composition of athletes in competitive sports and especially in youth competitive sports can also vary widely

within the same type of sport (34, 41). This becomes particularly clear in track and field, in which athletes in endurance sports weigh 50-60 kg and throwers can weigh up to 130 kg (33).

### Energy Requirements

While energy intake of humans is limited to their meals and possibly drinks, energy expenditure is a continuous process designed to maintain numerous activities and functions in our bodies. Daily energy requirements correspond to the energy expenditure over a 24-hour period (10). Accordingly, appropriate energy intake is a key factor in the human diet. Ideally, the energy intake matches the energy requirements. This is the amount of nutritional energy that guarantees a stable body weight and promotes health and fitness (10). However, energy requirement is not a fixed quantity and is influenced by numerous factors. These include e.g. body weight, body composition, physical activity, growth, pregnancy and nursing period. Physical activity is the most variable component in energy requirements (10, 44).

Adult endurance athletes report training programs ranging from 500 to 1,000 hours per year (17) or a running distance of >150 km per week (17, 36). In extreme situations such as the Tour de France or Race Across America, energy expenditure may be five times as much as when the athletes are at rest (18, 43). For individual long-distance races, this can result in an energy expenditure of >10,000 kcal per day (14). In contrast, there are also phases with low physical activity during a training year, in which daily energy requirements are at a lower level, owing to a potential training period of less than two hours per week. This makes it comparable to that of a person who exercises recreationally (17, 36, 37).

Based on the factors that influence the energy expenditure outlined above, the energy needed by most athletes lies between 1,500 kcal and 6,000 kcal per day and during the sports season can be estimated at 2,000 kcal to 5,000 kcal per day for a person weighing 70 kg, for example (37).

This variation in individual energy requirements must be considered during nutrition counseling. As a result, nutrition counseling professionals must address the type of sport/discipline and conditional training and competition exertions in order to get an accurate estimate of (current) energy requirements.

### Measurement and Estimate of Energy Requirements

Energy expenditure can be determined through various methods (e.g. direct calorimetry, indirect calorimetry, heart rate) (10). Indirect calorimetry is often used under lab conditions. In this process, the oxygen intake ( $\dot{V}O_2$ ) and carbon dioxide output ( $\dot{V}CO_2$ ) are measured via spirometry in liters per minute and calculated according to the Weir formula (42).

#### Weir Formula

$$\text{Energy (kcal/min)} = 3.9 \times \dot{V}O_2 \text{ (L/min)} + 1.1 \times \dot{V}CO_2 \text{ (L/min)}$$

Indirect calorimetry makes it possible to determine the resting energy expenditure, the basal metabolic rate and the total energy expenditure. Additionally,  $\dot{V}O_2$  and  $\dot{V}CO_2$  can be recorded with mobile devices during selected physical activities in the field. By using portable spirometry for a soccer training game,

it was possible to show an average energy consumption of 1,340 kcal for a period of 90 minutes, but there were great differences in the energy consumed by the studied players (1,050-1,750 kcal) (13). When measuring energy expenditure with spirometry, it should also be noted that the additional CO<sub>2</sub> formed by the bicarbonate buffer system with increased training intensity limits the accuracy of the values determined through indirect calorimetry for energy expenditure and substrate utilization. This applies particularly to physical exercise above a respiratory quotient (RQ) of 1.0.

Despite these limitations, it is possible to capture energy expenditure during physical activities through spirometric data for guidance. The results from numerous studies to determine the oxygen intake ( $\dot{V}O_2$ ) during selected physical activities were summarized in the Compendium of Physical Activities (2, 3, 4) and are updated regularly on the Compendium website (5). The Compendium has established itself as a widely used and accepted source to estimate energy consumption in recent decades (4, 39). The information in the Compendium is provided as a so-called "metabolic equivalent of task (MET)" (Box).

### Metabolic Equivalent of Task (MET)

One MET corresponds to resting energy expenditure and is defined as 1 kcal per kg of body weight per hour (kcal/kg body weight/hour). The MET values accordingly stand for a multiplication factor of resting energy expenditure (see examples 1 and 2). This is comparable with information about the Physical Activity Level (PAL). This is defined as the ratio of total energy expenditure to resting energy expenditure, but in contrast to MET, the applicable period is 24 hours. The PAL value thus refers to the additional daily energy consumption used during physical activities relative to the resting energy expenditure (10, 20).

The MET values describe specific activities, for which activity-related data are then available (Table 1). The MET values of the Compendium are in a range of 0.95 (sleeping) and 23 (running at 22.4 km/h) (4).

Based on the knowledge of resting energy expenditure (see formula for the calculation below), it is possible to calculate energy expenditure for different activities (Figure 1 with example 1 and 2).

This approach has established itself for estimating the energy expenditure during physical activity, but it comes with a few limitations that must be considered (4, 39). For example, the METs only apply for people without a mental or physical disability aged 18-65 years. It should also be considered that persons who are more physically fit reach a higher metabolic rate per unit of time with the same relative intensity of training (32). The intensity information (e.g. casual, general, strenuous) for the METs may be particularly misleading. Subjectively, a sports activity (e.g. soccer) can be perceived as very strenuous (e.g. owing to poor fitness). However, it is quite possible that a trained player has objectively run more and thus also shows greater energy expenditure while perceiving the activity "only" as moderately strenuous.

Additionally, a key factor in estimating energy expenditure is knowledge of resting energy expenditure. This can be determined with an indirect calorimetry, as described above. This is a complicated process, however, and is only rarely applied as a matter of routine. In general, the resting energy expenditure is thus calculated with available formulas.

#### Example 1:

##### Calculation of Energy Expenditure during Office Work

Adult with a body weight of 75 kg  
Resting energy expenditure (REE) approx. 1,800 kcal per day or 75 kcal per hour  
METs for "office work" (sitting tasks, light effort [e.g. office work...]) Code 11580) = 1.5  
REE per hour x METs = metabolic rate per hour  
75 kcal x 1.5 METs = 112.5 kcal per hour of office work

#### Example 2:

##### Calculation of Energy Expenditure during Soccer

Soccer player weighing 75 kg  
Resting energy expenditure (REE) approx. 1,800 kcal per day or 75 kcal per hour  
METs for "soccer competitive" (Code 15605) = 10  
REE per hour x METs = metabolic rate per hour  
75 kcal x 10 METs = 750 kcal per hour  
Corresponds to an estimated energy expenditure of 750 kcal per hour of soccer or 1,125 kcal for 90 minutes

Figure 1

Examples for calculation of energy expenditure for different activities.

### Conventional Formulas to Calculate Resting Energy Expenditure (REE)

#### Harris and Benedict (1918) (16)

Men: REE [kcal/day] = 66.5 + 13.8 x weight [kg] + 5.0 x size [cm] - 6.8 x age [years]  
Women: REE [kcal/day] = 655 + 9.6 x weight [kg] + 1.8 x size [cm] - 4.7 x age [years]

#### Cunningham (1980) (8)

REE [kcal/day] = 500 + 22 x LBM (Lean Body Mass)

#### Müller (2004) (10)

REE [MJ/day]\* = 0.047 x weight (kg) + 1.009 x gender (women = 0, men =1) - 0.01452 x age (years) + 3.21

\*Conversion of MJ into kcal by multiplying by 239

However, studies have shown that the calculated values do not always correspond to the measured values (7, 15). Although the mean values may generally coincide, there may still be a considerable discrepancy between measured and calculated resting energy expenditure in individual cases (7, 15). Since the fat-free body mass significantly influences the resting energy expenditure and is generally higher in athletes than non-athletes, it makes sense to use a corresponding calculation formula that takes fat-free body mass into account. The Cunningham formula (1980) is often applied in this respect. However, studies that compare the Cunningham formula with measured resting energy expenditure data show inconsistent results (7, 40).

Additionally, resting energy expenditure varies and may not be considered a fixed quantity. In some types of sports (e.g. weight-class sports, ski jumping, endurance sports), there are attempts to reduce body mass during selected periods in a season or to keep it low by means of a chronically low energy intake. This can lower resting energy expenditure during the season. However, this effect seems to be cancelled out at the end of the season with correspondingly higher energy intake (28). ➤

**Example 3:**

**Evaluation of Energy Availability**

Female athlete with a body weight of 60 kg, 20% body fat, 80% FFM (= 48.0 kg FFM)

Energy intake 2,600 kcal, energy expenditure during training 400 kcal

$$EA = 2,600 \text{ kcal} - 400 \text{ kcal} = 2,200 \text{ kcal}$$

$$2,200 \text{ kcal}/48 \text{ kg FFM} = 45.8 \text{ kcal/kg FFM}$$

**Example 4:**

**Energy Availability of a Female Athlete with Low Energy Intake and Increased Energy Expenditure During Training**

Female athlete with a body weight of 60 kg, 20% body fat, 80% FFM (= 48.0 kg FFM)

Energy intake 1,800 kcal, energy expenditure during training 600 kcal

$$EA = 1,800 \text{ kcal} - 600 \text{ kcal} = 1,200 \text{ kcal}$$

$$1,200 \text{ kcal}/48 \text{ kg FFM} = 25 \text{ kcal/kg FFM}$$

**Figure 2**

Examples for calculation of energy availability for different activities.

Particularly in competitive sports, it may not be possible to establish a universally valid formula to calculate resting energy expenditure. Reasons for this can be found in the large discrepancies in the area of the anthropometric data, phases of restrictive energy intake with an accompanying decrease in resting energy expenditure (28) as well as in the training phases that generally change, with varying training intensity and volume. Despite these described limitations, the use of formulas to calculate resting energy expenditure when supplementing the MET concept is considered useful for estimating energy requirements (39). However, it is always worth exercising sound judgment and care when evaluating the calculated result.

**Energy Availability**

In some types of sports or disciplines (e.g. ski jumping, high jump, marathon running), low body weight can provide an advantage in the performance or the regulations may necessitate a certain body weight on the day of the competition. To attain low body weight, athletes tend towards chronically low energy intake or, if possible, an increase in energy expenditure during training (24, 30). This is often referred to as a negative energy balance as well as low energy availability (EA).

**Energy Availability is Defined as:**

Energy availability = energy intake – energy expenditure during training

The EA is referred to in kcal per kg of fat-free mass (FFM) (Figure 2 with example 3 and 4).

The concept of energy availability is prevalent especially in sports nutrition and refers less to the traditional energy balance; instead it mainly considers the supplied amount of energy minus the energy used during sports. This results in the amount of energy available to the organism to maintain its basic functions (25, 39).

Studies involving female athletes show that low energy availability (<30 kcal per kg FFM) leads to a higher risk of signs of fatigue and overtraining, immunodeficiency, menstrual disorders and stress fractures (25, 26). Low energy availability is also seen as closely related to the development of the Female Athlete

Triad. This is a symptom complex related to menstrual cycle disorders, impaired eating behavior and lower bone density in female athletes (12, 31).

This phenomenon has not yet been systematically studied in men, but comparable results may be assumed (9, 21, 22). Since low energy availability can occur in both genders, it was suggested that the term Female Athlete Triad be replaced with the term Relative Energy Deficiency in Sports (RED-S) (29). This is also intended to take into account that low energy availability does not limit itself to bone metabolism and menstrual cycle disorders, but can also have hematological, metabolic, psychological, gastrointestinal, and immunological consequences. This is associated with negative influences on performance capacity, training adjustment, concentration, coordination, and an increased risk of injury (39). More in-depth information on this topic can be found in the freely accessible review by Nattiv et al. (2007) (31), De Souza et al. (2014) (12), Mountjoy et al. (2014) (29), Mountjoy et al. (2018) (30) and in a special issue of the International Journal of Sport Nutrition and Exercise Metabolism (Volume 28, Issue 4, July 2018).

Risk groups for low energy availability include athletes who deliberately ensure they have low body weight (e.g. weight reduction phase, permanently low body weight) or are pressured to maintain low body weight. Low energy availability may also affect athletes who e.g. do not eat enough for lack of time and/or underestimate their own energy requirements. It may also be attributable to the fact that hunger is suppressed for sports-related reasons, leading to insufficient energy intake (26). In phases of weight loss, it is important to ensure that energy availability of 30-45 kcal per kg FFM is guaranteed despite reduced energy intake (26).

When applying the concept of energy availability in a consulting practice, it must be kept in mind that information is needed on energy intake, energy expenditure during training as well as data regarding fat-free mass.

The limitations relating to determining energy expenditure are described above. General challenges in determining energy intake through the various nutrition survey systems are not part of this position paper but described in detail elsewhere (6, 27, 38).

Additionally, in nutrition surveys with athletes, it must be kept in mind that a one-time documentation of diets (e.g. once per year) will lead to a flawed evaluation of the nutritional situation, since energy intake can vary greatly owing to the different levels of exertion in training and competitions. The phenomena of underreporting, underrecording or underestimation and overestimation also occur in this group (6, 23, 27, 38). The significant causes named here include changes in eating behavior during the protocol phase, erroneous statements due to social desirability, and documentation errors in terms of the amount and description of the consumed foods (27). In nutrition surveys, athletes seem to “forget” to record such things as snacks or drinks, which can result in an erroneous evaluation of their energy intake (23, 27). Overall, this can lead to a misinterpretation of energy intake in the range of 10-45% (27) and in individual cases to a caloric deficit of >1,500 kcal per day (7).

Additionally, the body composition must be measured to obtain information on fat-free mass. Different results may be obtained depending on the measuring method (1), however, which, in turn, influences the calculation of energy availability.

Despite the limitations described above, it is recommended that more attention be paid to the topic of energy requirements in sports so that warning signals for low energy availability and the associated negative consequences on health and performance can be detected at an early stage.

## Conclusion

Depending on the training and competition phase within the yearly cycle, there can be enormous differences in energy requirements of athletes. Determining energy expenditure precisely is not easy, but can nevertheless be approximated. Energy intake that is adapted to individual needs is a key cornerstone in sports nutrition. In general, it is thus important to ensure adequate energy intake appropriate for the situation in order to meet the training and potentially growth-related needs. Inadequate energy intake can lead to low energy availability with the previously described consequences for health and performance. ■

## Acknowledgement

The authors wish to thank Dr. Angela Bechthold and Birte Petersen-Sperlich from the Science Department at the DGE.

## Conflict of Interest

*The authors have no conflict of interest.*

## References

- (1) ACKLAND TR, LOHMAN TG, SUNDGOT-BORGEN J, MAUGHAN JM, MEYER NL, STEWART AD, MÜLLER W. Current Status of Body Composition Assessment in Sport. Review and Position Statement on Behalf of the Ad Hoc Research Working Group on Body Composition Health and Performance, Under the Auspices of the I.O.C. Medical Commission. *Sports Med.* 2012; 42: 227-249. doi:10.2165/11597140-000000000-00000
- (2) AINSWORTH BE, HASKELL WL, LEON AS, JACOBS DR, MONTROYE HJ, SALLIS JF, PAFFENBARGER RS. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc.* 1993; 25: 71-80. doi:10.1249/00005768-199301000-00011
- (3) AINSWORTH BE, HASKELL WL, WHITT MC, IRWIN ML, SWARTZ AM, STRATH SJ, O'BRIEN WL, BASSETT DR, SCHMITZ KH, EMPLAINCOURT PO, JACOBS DR, LEON AS. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc.* 2000; 32: S498-S516. doi:10.1097/00005768-200009001-00009
- (4) AINSWORTH BE, HASKELL WL, HERRMANN SD, MECKES N, BASSETT DR, TUDOR-LOCKE C, GREER JL, VEZINA J, WHITT-GLOVER MC, LEON AS. 2011 Compendium of physical activities: a second update of codes and MET values. *Med Sci Sports Exerc.* 2011; 43: 1575-1581. doi:10.1249/MSS.0b013e31821ece12
- (5) AINSWORTH BE, HASKELL WL, HERRMANN SD, MECKES N, BASSETT JR DR, TUDOR-LOCKE C, GREER JL, VEZINA J, WHITT-GLOVER MC, LEON AS. The Compendium of Physical Activities Tracking Guide. Healthy Lifestyles Research Center, College of Nursing & Health Innovation, Arizona State University. <https://sites.google.com/site/compendiumofphysicalactivities/> [July 01<sup>st</sup> 2020].
- (6) BLACK AE. Dietary assessment for sports dietetics. *Nutr Bull.* 2001; 26: 29-42. doi:10.1046/j.1467-3010.2001.00096.x
- (7) CARLSOHN A, ROSENBERGER F, SCHAPP L, FUSCH G, MAYER F. Validität der Energiezufuhrbestimmung mittels Ernährungsprotokoll bei Normalgewichtigen in Abhängigkeit von der Höhe der Energiezufuhr. *Ernähr-Umsch.* 2012; 59: 572-577. doi:10.4455/ eu.2012.958
- (8) CUNNINGHAM JJ. A reanalysis of the factor influencing basal metabolic rate in normal adults. *Am J Clin Nutr.* 1980; 33: 2372-2374. doi:10.1093/ajcn/33.11.2372
- (9) DEGOUTTE F, OUANEL P, BÈGUE RJ, COLOMBIER M, LAC G, PEQUIGNOT JM, FILAIRE E. Food restriction, performance, biochemical, psychological, and endocrine changes in judo athletes. *Int J Sports Med.* 2006; 27: 9-18. doi:10.1055/s-2005-837505
- (10) DEUTSCHE GESELLSCHAFT FÜR ERNÄHRUNG, ÖSTERREICHISCHE GESELLSCHAFT FÜR ERNÄHRUNG, SCHWEIZERISCHE GESELLSCHAFT FÜR ERNÄHRUNG (HRSG.): Energie. In: Referenzwerte für die Nährstoffzufuhr. Bonn 2. Auflage. 1. Ausgabe, 2015, 1-20.
- (11) DESBROW B, MCCORMACK J, BURKE LM, COX GR, FALLON K, HISLOP M, LOGAN R, MARINO N, SAWYER SM, SHAW G, STAR A, VIDGEN H, LEVERITT M. Sports Dietitians Australia position statement: sports nutrition for the adolescent athlete. *Int J Sport Nutr Exerc Metab.* 2014; 24: 570-584. doi:10.1123/ijsnem.2014-0031
- (12) DE SOUZA MJ, NATTIV A, JOY E, MISRA M, WILLIAMS NI, MALLINSON RJ, GIBBS JC, OLMSTED M, GOOLSBY M, MATHESON G. 2014 Female Athlete Triad Coalition Consensus Statement on Treatment and Return to Play of the Female Athlete Triad: 1st International Conference held in San Francisco, California, May 2012 and 2nd International Conference held in Indianapolis, Indiana, May 2013. *Br J Sports Med.* 2014; 48: 289. doi:10.1136/bjsports-2013-093218
- (13) FERRAUTI A, GIESEN HT, MERHEIM G, WEBER K. Indirekte Kalorimetrie im Fußballspiel. *Dtsch Z Sportmed.* 2006; 57: 142-146.
- (14) GEESMANN B, MESTER J, KOEHLER K. Energy balance, macronutrient intake, and hydration status during a 1,230 km ultra-endurance bike marathon. *Int J Sport Nutr Exerc Metab.* 2014; 24: 497-506. doi:10.1123/ijsnem.2013-0169
- (15) HASSON RE, HOWE CA, JONES BL, FREEDSON PS. Accuracy of four resting metabolic rate prediction equations: effects of sex, body mass index, age, and race/ethnicity. *J Sci Med Sport.* 2011; 14: 344-351. doi:10.1016/j.jsams.2011.02.010
- (16) HARRIS JA, BENEDICT FG. A biometric study of basal metabolism in man. *Proc Natl Acad Sci USA.* 1918; 4: 370-373. doi:10.1073/pnas.4.12.370
- (17) HEYDENREICH J, KAYSER B, SCHUTZ Y, MELZER K. Total Energy Expenditure, Energy Intake, and Body Composition in Endurance Athletes Across the Training Season: A Systematic Review. *Sports Med Open.* 2017; 3: 8. doi:10.1186/s40798-017-0076-1
- (18) HULTON AT, LAHART I, WILLIAMS KL, GODFREY R, CHARLESWORTH S, WILSON M, PEDLAR C, WHYTE G. Energy expenditure in the Race Across America (RAAM). *Int J Sports Med.* 2010; 31: 463-467. doi:10.1055/s-0030-1251992

- (19) MAUGHAN RJ, SHIRREFFS SM. IOC Consensus Conference on Nutrition in Sport, 25–27 October 2010, International Olympic Committee, Lausanne, Switzerland. *J Sports Sci.* 2010; 29: S1. doi:10.1080/02640414.2011.619339
- (20) INSTITUTE OF MEDICINE. Dietary Reference Intakes: The Essential Guide to Nutrient Requirements. Washington, DC: The National Academies Press. 2006. doi:10.17226/11537
- (21) KARILA TAM, SARKKINEN P, MARTTINEN M, SEPPÄLÄ T, MERO A, TALLROTH K. Rapid weight loss decreases serum testosterone. *Int J Sports Med.* 2008; 29: 872–877. doi:10.1055/s-2008-1038604
- (22) KOEHLER K, ACHTZEHN S, BRAUN H, MESTER J, SCHAEZNER W. Comparison of self-reported energy availability and metabolic hormones to assess adequacy of dietary energy intake in young elite athletes. *Appl Physiol Nutr Metab.* 2013; 38: 725–733. doi:10.1139/apnm-2012-0373
- (23) LARSON-MEYER DE, WOOLF K, BURKE L. Assessment of Nutrient Status in Athletes and the Need for Supplementation. *Int J Sport Nutr Exerc Metab.* 2018; 28: 139–158. doi:10.1123/ijsnem.2017-0338
- (24) LOGUE D, MADIGAN SM, DELAHUNT E. Low Energy Availability in Athletes: A Review of Prevalence, Dietary Patterns, Physiological Health, and Sports Performance. *Sports Med.* 2018; 48: 73–96. doi:10.1007/s40279-017-0790-3
- (25) LOUCKS AB. Energy balance and body composition in sports and exercise. *J Sports Sci.* 2004; 22: 1–14. doi:10.1080/0264041031000140518
- (26) LOUCKS AB, KIENS B, WRIGHT HH. Energy availability in athletes. *J Sports Sci.* 2011; 29: S7–S15. doi:10.1080/02640414.2011.588958
- (27) MAGKOS F, YANNAKOULIA M. Methodology of dietary assessment in athletes: concepts and pitfalls. *Curr Opin Clin Nutr Metab Care.* 2003; 6: 539–549. doi:10.1097/00075197-200309000-00007
- (28) MELBY CL, SCHMIDT WD, CORRIGAN D. Resting metabolic rate in weightcycling collegiate wrestlers compared with physically active, noncycling control subjects. *Am J Clin Nutr.* 1990; 52: 409–414. doi:10.1093/ajcn/52.3.409
- (29) MOUNTJOY M, SUNDGOT-BORGEN J, BURKE L, CARTER S, CONSTANTINI N, LEBRUN C, MEYER N, SHERMAN R, STEFFEN K, BUDGETT R, LJUNGQVIST A. The IOC consensus statement: beyond the Female Athlete Triad—Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med.* 2014; 48: 491–497. doi:10.1136/bjsports-2014-093502
- (30) MOUNTJOY M, SUNDGOT-BORGEN JK, BURKE LM, ACKERMAN KE, BLAUWET C, CONSTANTINI N, LEBRUN C, LUNDY B, MELIN AK, MEYER N, SHERMAN RT, TENFORDE AS, TORSTVEIT MK, BUDGETT R. IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med.* 2018; 52: 687–697. doi:10.1136/bjsports-2018-099193
- (31) NATTIV A, LOUCKS AB, MANORE MM, SANBORN CF, SUNDGOT-BORGEN J, WARREN MP. American College of Sports Medicine position stand. The female athlete triad. *Med Sci Sports Exerc.* 2007; 39: 1867–1882.
- (32) NORTON K, NORTON L, SADGROVE D. Position statement on physical activity and exercise intensity terminology. *J Sci Med Sport.* 2010; 13: 496–502. doi:10.1016/j.jsams.2009.09.008
- (33) O'CONNOR H, OLDS T, MAUGHAN RJ. Physique and performance for track and field events. *J Sports Sci.* 2007; 25: S49–S60. doi:10.1080/02640410701607296
- (34) SANTOS DA, DAWSON JA, MATIAS CN, ROCHA PM, MINDERICO CS, ALLISON DB, SARDINHA LB, SILVA AM. Reference Values for Body Composition and Anthropometric Measurements in Athletes. *PLoS One.* 2014; 9: e97846. doi:10.1371/journal.pone.0097846
- (35) SPOMEDIAL (HRSG.). Der respiratorische Quotient (RQ). [http://vmrz0100.vmr.uni-bochum.de/spomedial/content/e866/e2442/e3862/e4009/e4038/index\\_ger.html](http://vmrz0100.vmr.uni-bochum.de/spomedial/content/e866/e2442/e3862/e4009/e4038/index_ger.html) [July 01<sup>st</sup> 2020]
- (36) STELLINGWERFF T, BOIT MK, RES PT. Nutritional strategies to optimize training and racing in middle-distance athletes. *J Sports Sci.* 2007; 25: S17–S28. doi:10.1080/02640410701607213
- (37) STELLINGWERFF T, MAUGHAN RM, BURKE LM. Nutrition for power sports: Middle-distance running, track cycling, rowing, canoeing/kayaking, and swimming. *J Sports Sci.* 2011; 29: S79–S89. doi:10.1080/02640414.2011.589469
- (38) STRASSBURG A. Ernährungserhebungen – Methoden und Instrumente. *Ernahr-Umsch.* 2010; 10: 422–430.
- (39) THOMAS DT, ERDMAN KA, BURKE LM. American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance. *Med Sci Sports Exerc.* 2016; 48: 543–568.
- (40) THOMPSON JL, MANORE MM. Predicted and measured resting metabolic rate of male and female endurance athletes. *J Am Diet Assoc.* 1996; 96: 30–34. doi:10.1016/S0002-8223(96)00010-7
- (41) VAN ERP-BAART AM, SARIS WH, BINKHORST RA, VOS JA, ELVERS JW. Nationwide survey on nutritional habits in elite athletes. Part I. Energy, carbohydrate, protein, and fat intake. *Int J Sports Med.* 1989; 10: S3–S10. doi:10.1055/s-2007-1024947
- (42) WEIR JB. New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol.* 1949; 109: 1–9. doi:10.1113/jphysiol.1949.sp004363
- (43) WESTERTERP KR, SARIS WH, VAN ES M, TEN HOOR F. Use of the doubly labeled water technique in humans during heavy sustained exercise. *J Appl Physiol.* 1986; 61: 2162–2167. doi:10.1152/jappl.1986.61.6.2162
- (44) WESTERTERP KR. Daily physical activity as determined by age, body mass and energy balance. *Eur J Appl Physiol.* 2015; 115: 1177–1184. doi:10.1007/s00421-015-3135-7